



GABRA1 gene

gamma-aminobutyric acid type A receptor alpha1 subunit

Normal Function

The *GABRA1* gene provides instructions for making one piece, the alpha-1 ($\alpha 1$) subunit, of the GABA_A receptor protein. GABA_A receptors are made up of different combinations of five protein subunits, each produced from a different gene. (Nineteen different genes provide instructions for GABA_A receptor subunits.) These subunits form a hole (pore) in the cell membrane through which negatively charged chlorine atoms (chloride ions) can flow.

A chemical that transmits signals in the brain (a neurotransmitter) called gamma-amino butyric acid (GABA) attaches to GABA_A receptors. Once GABA attaches, the pore formed by the subunits opens, and chloride ions flow across the cell membrane. After infancy, chloride ions flow into the cell through the open pore, which creates an environment in the cell that blocks (inhibits) signaling between neurons. The primary role of GABA in children and adults is to prevent the brain from being overloaded with too many signals. In contrast, in newborns and infants, chloride ions flow out of the cell when the pore is opened, creating an environment that allows signaling between neurons.

Health Conditions Related to Genetic Changes

juvenile myoclonic epilepsy

A mutation in the *GABRA1* gene has been identified in at least one family with juvenile myoclonic epilepsy. This condition typically begins in childhood or adolescence and causes recurrent myoclonic seizures, which are characterized by rapid, uncontrolled muscle jerks. Affected individuals can also have other types of seizures called generalized tonic-clonic seizures (or grand mal seizures) and absence seizures. The mutation associated with this condition changes a single protein building block (amino acid) in the $\alpha 1$ subunit. The amino acid alanine at protein position 322 is replaced by the amino acid asparagine. This gene mutation is written as Ala322Asp or A322D.


This *GABRA1* gene mutation leads to the formation of an abnormal $\alpha 1$ subunit that reduces GABA_A receptor function. GABA_A receptors containing the abnormal subunit are broken down before they reach the cell membrane. Studies show that the altered receptors can also interfere with normal receptors inside the cell, leading to the additional loss of normal receptors. Because of the reduction of GABA_A

receptor function, signaling between neurons is not regulated, which can lead to overstimulation of neurons. Researchers believe that the overstimulation of certain neurons in the brain triggers the abnormal brain activity associated with seizures.

Chromosomal Location

Cytogenetic Location: 5q34, which is the long (q) arm of chromosome 5 at position 34

Molecular Location: base pairs 161,847,191 to 161,899,959 on chromosome 5 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- ECA4
- EJM
- EJM5
- GABA(A) receptor subunit alpha-1
- GABA(A) receptor, alpha 1
- gamma-aminobutyric acid (GABA) A receptor, alpha 1
- gamma-aminobutyric acid receptor subunit alpha-1
- gamma-aminobutyric acid receptor subunit alpha-1 precursor
- GBRA1_HUMAN

Additional Information & Resources

Educational Resources

- Neuroscience (second edition, 2001): GABA and Glycine
<https://www.ncbi.nlm.nih.gov/books/NBK11084/>
- Neuroscience (second edition, 2001): GABA and Glycine Receptors
<https://www.ncbi.nlm.nih.gov/books/NBK10977/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28GABRA1%5BTIAB%5D%29+OR+%28%28EJM%5BTIAB%5D%29+OR+%28ECA4%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2160+days%22%5Bdp%5D>

OMIM

- GAMMA-AMINOBUTYRIC ACID RECEPTOR, ALPHA-1
<http://omim.org/entry/137160>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_GABRA1.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=GABRA1%5Bgene%5D>
- HGNC Gene Family: Gamma-aminobutyric acid type A receptor subunits
<http://www.genenames.org/cgi-bin/genefamilies/set/563>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=4075
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/2554>
- UniProt
<http://www.uniprot.org/uniprot/P14867>

Sources for This Summary

- Ben-Ari Y, Khalilov I, Kahle KT, Cherubini E. The GABA excitatory/inhibitory shift in brain maturation and neurological disorders. *Neuroscientist*. 2012 Oct;18(5):467-86. Epub 2012 Apr 30. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22547529>
- Bradley CA, Taghibiglou C, Collingridge GL, Wang YT. Mechanisms involved in the reduction of GABAA receptor alpha1-subunit expression caused by the epilepsy mutation A322D in the trafficking-competent receptor. *J Biol Chem*. 2008 Aug 8;283(32):22043-50. doi: 10.1074/jbc.M801708200. Epub 2008 Jun 5.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18534981>
- Cossette P, Liu L, Brisebois K, Dong H, Lortie A, Vanasse M, Saint-Hilaire JM, Carmant L, Verner A, Lu WY, Wang YT, Rouleau GA. Mutation of GABRA1 in an autosomal dominant form of juvenile myoclonic epilepsy. *Nat Genet*. 2002 Jun;31(2):184-9. Epub 2002 May 6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11992121>

- Ding L, Feng HJ, Macdonald RL, Botzolakis EJ, Hu N, Gallagher MJ. GABA(A) receptor alpha1 subunit mutation A322D associated with autosomal dominant juvenile myoclonic epilepsy reduces the expression and alters the composition of wild type GABA(A) receptors. *J Biol Chem*. 2010 Aug 20;285(34):26390-405. doi: 10.1074/jbc.M110.142299. Epub 2010 Jun 15.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20551311>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2924069/>
- OMIM: GAMMA-AMINOBUTYRIC ACID RECEPTOR, ALPHA-1
<http://omim.org/entry/137160>
- Gallagher MJ, Ding L, Maheshwari A, Macdonald RL. The GABAA receptor alpha1 subunit epilepsy mutation A322D inhibits transmembrane helix formation and causes proteasomal degradation. *Proc Natl Acad Sci U S A*. 2007 Aug 7;104(32):12999-3004. Epub 2007 Aug 1.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17670950>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1941799/>
- Kang JQ, Shen W, Macdonald RL. Two molecular pathways (NMD and ERAD) contribute to a genetic epilepsy associated with the GABA(A) receptor GABRA1 PTC mutation, 975delC, S326fs328X. *J Neurosci*. 2009 Mar 4;29(9):2833-44. doi: 10.1523/JNEUROSCI.4512-08.2009.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19261879>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2687144/>
- Krampfl K, Maljevic S, Cossette P, Ziegler E, Rouleau GA, Lerche H, Bufler J. Molecular analysis of the A322D mutation in the GABA receptor alpha-subunit causing juvenile myoclonic epilepsy. *Eur J Neurosci*. 2005 Jul;22(1):10-20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16029191>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/GABRA1>

Reviewed: June 2012
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services